

breakfast

Birth through 5 months	6 through 11 months
4-6 fluid ounces breastmilk ¹ or formula ²	6-8 fluid ounces breastmilk ¹ or formula ² ; and 0 to ½ ounce equivalent (0-4 tablespoons) infant cereal ^{2,3} or, 0-4 tablespoons meat, fish, poultry, whole egg, cooked dry beans, or cooked dry peas; or 0-2 ounces of cheese; or 0-4 ounces (volume) of cottage cheese; or 0-4 ounces or ½ cup of yogurt ⁴ ; or a combination of the above ⁵ ; and 0-2 tablespoons vegetable or fruit or a combination of both ^{5,6}

¹ Breastmilk or formula, or portions of both, must be served; however, it is recommended that breastmilk be served in place of formula from birth through 11 months. For some breastfed infants who regularly consume less than the minimum amount of breastmilk per feeding, a serving of less than the minimum amount of breastmilk may be offered, with additional breastmilk offered at a later time if the infant will consume more.

² Infant formula and dry infant cereal must be iron-fortified.

³ Ounce equivalents (oz eq) are used to determine the quantity of creditable grains.

⁴ Yogurt must contain no more than 23 grams of total sugars per 6 ounces.

⁵ A serving of this component is required when the infant is developmentally ready to accept it.

⁶ Fruit and vegetable juices must not be served.

lunch and supper

Birth through 5 months	6 through 11 months
4-6 fluid ounces breastmilk ¹ or formula ²	6-8 fluid ounces breastmilk ¹ or formula ² ; and 0 to ½ ounce equivalent (0-4 tablespoons) infant cereal ^{2,3} or, 0-4 tablespoons meat, fish, poultry, whole egg, cooked dry beans, or cooked dry peas; or 0-2 ounces of cheese; or 0-4 ounces (volume) of cottage cheese; or 0-4 ounces or ½ cup of yogurt ⁴ ; or a combination of the above ⁵ ; and 0-2 tablespoons vegetable or fruit or a combination of both ^{5,6}

¹ Breastmilk or formula, or portions of both, must be served; however, it is recommended that breastmilk be served in place of formula from birth through 11 months. For some breastfed infants who regularly consume less than the minimum amount of breastmilk per feeding, a serving of less than the minimum amount of breastmilk may be offered, with additional breastmilk offered at a later time if the infant will consume more.

² Infant formula and dry infant cereal must be iron-fortified.

³ Ounce equivalents (oz eq) are used to determine the quantity of creditable grains.

⁴ Yogurt must contain no more than 23 grams of total sugars per 6 ounces.

⁵ A serving of this component is required when the infant is developmentally ready to accept it.

⁶ Fruit and vegetable juices must not be served.

snack

Birth through 5 months	6 through 11 months
4-6 fluid ounces breastmilk ¹ or formula ²	2-4 fluid ounces breastmilk ¹ or formula ² ; and
	0-½ oz eq bread or bread-like items ^{3,4} ; or 0-¼ oz eq crackers ^{3,4} ; or 0-½ oz eq (4 tablespoons) infant cereal ^{2,3,4} or 0-¼ oz eq ready-to-eat breakfast cereal ^{3,4,5,6} ; and
	0-2 tablespoons vegetable or fruit or a combination of both ^{6,7}

¹ Breastmilk or formula, or portions of both, must be served; however, it is recommended that breastmilk be served in place of formula from birth through 11 months. For some breastfed infants who regularly consume less than the minimum amount of breastmilk per feeding, a serving of less than the minimum amount of breastmilk may be offered, with additional breastmilk offered at a later time if the infant will consume more.

² Infant formula and dry infant cereal must be iron-fortified.

³ Ounce equivalents (oz eq) are used to determine the quantity of creditable grains.

⁴ A serving of grains must be whole grain-rich, enriched meal, or enriched flour.

⁵ Breakfast cereals must contain no more than 6 grams of sugar per dry ounce (no more than 21.2 grams sucrose and other sugars per 100 grams of dry cereal).

⁶ A serving of this component is required when the infant is developmentally ready to accept it.

⁷ Fruit and vegetable juices must not be served.

Infant Meal Pattern

This project has been funded at least in part with Federal funds from the U.S. Department of Agriculture, Food and Nutrition Service through an agreement with the Institute of Child Nutrition at the University of Mississippi. The contents of this publication do not necessarily reflect the views or policies of the U.S. Department of Agriculture, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. government.

The University of Mississippi is an EEO/AA/TitleVI/Title IX/Section 504/ADA/ADEA Employer.

In accordance with Federal law and U.S. Department of Agriculture policy, this institution is prohibited from discriminating on the basis of race, color, national origin, sex, age, or disability. To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 1400 Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

© 2022, Institute of Child Nutrition, The University of Mississippi, School of Applied Sciences

Except as provided below, you may freely use the text and information contained in this document for non-profit or educational use with no cost to the participant for the training providing the following credit is included. These materials may not be incorporated into other websites or textbooks and may not be sold. The photographs and images in this document may be owned by third parties and used by the University of Mississippi under a licensing agreement. The University cannot, therefore, grant permission to use these images.

For more information, please contact helpdesk@theicn.org.

07/07/2022